arXiv:1304.2197v1 [quant-ph] 8 Apr 2013

An extension of the Wigner inequality: theory and experiment

William N. Plick,[1,2] Robert Fickler,[1,2] Radek Lapkiewicz,[1,2] and Sven Ramelow[1,2]

[1]*Quantum Optics, Quantum Nanophysics, Quantum Information,*
University of Vienna, Boltzmanngasse 5, Vienna A-1090, Austria
[2]*Institute for Quantum Optics and Quantum Information, Boltzmanngasse 3, Vienna A-1090, Austria*

(Dated: April 9, 2013)

In the study of systems that can not be explained classically, the Wigner inequality, has received only a small amount of attention. In this paper we extend the Wigner inequality – originally derived in 1969 – and show how it may be used to contradict local realism with only coincidence detections in the absence of two-outcome measurements – that is, any system where only one possible result of a pair of potential outcomes can be registered. This encapsulates a much broader class of measurement schemes than could previously violate a local-realistic inequality. We then apply this inequality to a recently constructed setup with access to entangled pairs of photons with very high angular momenta, in which no previously derived local-realistic inequality could successfully be used without making very broad assumptions. This experiment constitutes the first violation of local realism with such high quanta. We thus demonstrate the versatility of this inequality under very lossy conditions.

PACS numbers: 03.67.Hk, 03.65.Ud, 42.50.Xa, 42.50.-p

The ability to banish the local-realistic world-view can be an important benchmark for experiments in quantum physics. Much effort has been dedicated in recent decades to highlight the contradictory predictions of classical and quantum mechanics with ever larger and less conventional systems. As a result, in recent years the number and complexity of experiments whose results can not be explained using local-realistic theories has increased dramatically. At the forefront of those efforts are the experiments which seek to entangle novel quantum systems. Many inequalities exist whose violation provides evidence against the proposition of local-realism (under reasonable assumptions) and thus shows quantum behavior.

These inequalities require either dichotomic (that is, two-outcome) measurements on correlations (coincidence counts, photonically), or the use of local measurements (singles, photonically) [1–3]. Examples of dichotomic measurements are spin up or spin down, horizontal or vertical polarization, etc. Implementing the two-outcome measurements required by most tests of classical-realism can be difficult. The other option is to explicitly include the single count rates, which typically are much higher than coincidences, requiring very high efficiency. For example, in the seminal paper by Aspect [4] only polarizers are used – not polarizing beam splitters – and in fact it is not until later [5] that true dichotomic measurements are used, allowing violation without reconstructing the statistics for dichotomic outcomes based on the statistics for single-outcome measurements.

An exception to these conditions is the Wigner Inequality (WI), which needs only single-outcome coincidences [6]. The price for this flexibility is that the original derivation of this inequality assumes perfect anti-correlations.

Stated most concisely, in its simplest form as derived in 1969, the Wigner inequality gives a restriction on the statistical measurement outcomes of potential hidden variable models (HVM) of perfectly anti-correlated, two-particle, spin-1/2 systems – given three potential measurement settings. Specifically, it states that the probabilities of getting coincident detections (either spin-up, spin-up; or spin-down, spin-down) must obey the relation $P_{13} \leq P_{12} + P_{23}$ where the subscripts indicate one of the three settings (typically projection angles) on either the first or second particle (indicated by first or second position in the subscript). This inequality can be violated quantum mechanically with a singlet state by $3/4 \leq 1/4 + 1/4$. These basic statements will be derived carefully and expanded upon in what follows.

Modified versions of the WI have been found which do not require perfect anti-correlations, however these again require dichotomic measurements [7, 8]. In this paper we derive an extended version of the Wigner inequality which does not require dichotomic measurements, use of the singles, or perfect anti-correlation. This allows violation of local realism in a much broader class of experimental systems than was possible before.

In the following section an extended version of the original Wigner inequality is derived and we carefully outline the necessary assumptions, including how we maintain the single-outcome-only measurements despite the reality of imperfect anti-correlations. In the third section we apply the newly derived inequality to a recent experiment [11] utilizing very high orbital angular momentum – a situation in which other tests of local realism may not be applied (due to the nature of the detection scheme). In the final section we summarize the main points of the paper.

Derivation of the General Inequality– Following Wigner [6], assume that there exists a deterministic description of the world which relies on a hidden variable – which complectly defines the results of all possible measurements, but which is not in principle known or knowable. When a measurement is performed, the statistical distribution of the hidden variable reproduces the probabilities for the potential outcomes. We need not be concerned about the actual statistical distribution of the variable, but only define domains of it which correspond to obtaining a given result when a particular measurement is performed. Each of these domains will be represented by a symbol which, herein, will be called a “Wigner-symbol”.

We will take the case of a two particle system, where some degree of anti-correlation is present, and assume that these anti-correlations can be explained via classical statistics. We limit the measurements settings to three possibilities for each particle ϕ_1, ϕ_2, and ϕ_3. We limit the measurement outcomes to two possibilities: a detection, or no detection. So, we can define the Wigner-symbols (representing domains of the space of the statistical distribution of the hidden variable) in the form $(\phi_{1a}, \phi_{2a}, \phi_{3a}; \phi_{1b}, \phi_{2b}, \phi_{3b})$, where the ϕ's can take the values of “+” or “−”, indicating a detection, or no detection, respectively at the setting, and particle (a or b), indicated by the subscripts. If we had the case of perfect anti-correlation we could write only symbols which exhibited this quality, like $(++-,--+)$. This symbol would mean, for example, if a ϕ_1 measurement were performed on particle a and a ϕ_2 measurement were performed on particle b the results would be a detection and no detection, respectively. However, we must allow for the possibility of symbols which do not represent perfect anti-correlations, as particle pairs which are not perfectly anti-correlated can and do occur. This generality is an extension to the original inequality as derived by Wigner.

The Wigner-symbols also have the direct interpretation as the probabilities that the hidden variables take a value lying in the domain specified by that symbol. That is, the size of the domain corresponding to the indicated measurement outcomes is the probability of such an outcome occurring. Note, that we have made no assumptions about how the symbols are weighted and thus have made no assumptions about the statistical distribution of the hidden variables. Now, if we are to assume that there exists a realistic description of the world, the probabilities for some chosen outcomes of the described setup can be written as

$$
\begin{aligned}
P_{13} &= (++-,--+) + (+--,-++) + \mathcal{S}_{13}, \quad (1)\\
P_{12} &= (+-+,-+-) + (+--,-++) + \mathcal{S}_{12}, \quad (2)\\
P_{23} &= (-+-,+-+) + (++-,--+) + \mathcal{S}_{23}. \quad (3)
\end{aligned}
$$

Where the P_{ij}'s are the probabilities of getting a coincident detection when the detectors are set to the measurement positions indicated by the subscripts. For example, P_{13} is the probability that we detect particles a and b if the measuring devices are in settings ϕ_1 and ϕ_3 respectively. The symbols $\mathcal{S}_{ij}$ represent a collection of those Wigner-symbols which do not exhibit the perfect anti-correlation, for example $(+++,--+)$. There are 14 such Wigner-symbols included in each $\mathcal{S}_{ij}$. Putting Eqs.(2) and (3) in a slightly different form

$$
\begin{aligned}
P_{12} - (+-+,-+-) - \mathcal{S}_{12} &= (+--,-++), \quad (4)\\
P_{23} - (-+-,+-+) - \mathcal{S}_{23} &= (++-,--+). \quad (5)
\end{aligned}
$$

And then using this, we write

$$
\begin{aligned}
P_{12} - \mathcal{S}'_{12} &\geq P_{12} - (+-+,-+-) - \mathcal{S}_{12},\\
&\geq (+--,-++), \quad (6)\\
P_{23} - \mathcal{S}'_{23} &\geq P_{23} - (-+-,+-+) - \mathcal{S}_{23},\\
&\geq (++-,--+). \quad (7)
\end{aligned}
$$

Where $\mathcal{S}'_{12}$ and $\mathcal{S}'_{23}$ consist of only those Wigner-symbols representing imperfect anti-correlations for which the same symbol exists in $\mathcal{S}_{13}$. Direct substitution of Eq.(6) and Eq.(7) into Eq.(1) yields $P_{13} - \mathcal{S} \leq P_{12} + P_{23}$. Where $\mathcal{S}$ collects all those Wigner-symbols which exhibit imperfect anti-correlations which can not be canceled out. If we were to take $\mathcal{S}$ to zero we would recover the original result by Wigner. Now,

$$
\begin{aligned}
\mathcal{S} &= \mathcal{S}_{13} - \mathcal{S}'_{12} - \mathcal{S}'_{23}\\
&= (+--,--+) + (+-+,--+)\\
&\quad +(+--,+-+) + (+-+,+-+). \quad (8)
\end{aligned}
$$

The first line explicitly includes all the symbols lacking from Wigner's original derivation. There are 28 terms in total, of which 24 cancel out. The fact that these cancelations can be made is a direct result of the assumption of the independence of measurement context. In the second line we write the result. The second and third terms can be bounded from above by observing the coincident intensity at a measurement setting where there is a minimum – at ϕ_{a3} and ϕ_{b3} for the second term, and ϕ_{a1} and ϕ_{b1} for the third. Either of these measurements will also set an upper-bound on the fourth term. More problematic is the first term, if dichotomic measurements were available an upper-bound could be placed on this term directly, by making measurements of “no-click” outcomes. However, in their absence, this term will be estimated via an extended assumption of fairness.

The assumption of fairness of measurement results relates to a loophole in Bell and CHSH-type inequalities, known formally as “fair sampling”. This loophole can be closed by increasing the detector efficiency to a suitable degree [9]. Though any particular application of the extended Wigner inequality may, or may not, be susceptible to a fair sampling loophole our assumption of “extended fairness” differs. It is however philosophically reminiscent, in the sense that it is in some way an assumption that the hidden variable model is not “conspiring”.

Precisely, our assumption is that: An imperfect correlation is not more likely to occur when detectors are set to any particular measurement position. This is a less well-founded assumption than fair sampling – as fair sampling assumes that the HVM allows the detection device to fairly sample the ensemble – whereas extended fairness assumes that the detection device, and whatever object allows projections in the Hilbert space (eg. polarizer) are both acting fairly and not influencing or influenced by the HVM.

It is important to point out that – while our assumption is a somewhat broader one than fair sampling due to the fact that is physically includes more of the experimental setup – it is still a more well founded assumption than the "no-enhancement assumption" of Clauser and Horne [10]. The no-enhancement assumption states that the addition of a polarizer can not enhance the probability of a detection. Our assumption allows for the polarizer (or other device) to enhance, degrade, or have no effect on counts. Furthermore a Bell-type inequality derived using no-enhancement requires correlation measurements with the polarizers (or other devices) removed, which can create problems in implementation (physical difficulty, and skewed count rates).

With the addition of this assumption we can take the first term of Eq.(8) and determine which symbols representing perfect correlations could give rise to the $\mathcal{S}$ term of interest under some physical process (dark count, scattering, etc.). Both $(+--,-++)$ and $(++-,--+)$ could give rise to $(+--,--+)$ after a single physical event. For each of these symbols representing a perfect correlation there are six symbols which represent a single event occurring to that symbol. The fundamental application of our assumption is that each of these twelve symbols are equally likely, and that therefore we may replace one with another without changing the probabilities of the outcome. We take $(+--,--+)$ and replace it with $(+--,+++)$. Similarly we can replace $(+-+,--+)$ with $(+--,++-)$ (this second replacement is not strictly necessary, though it simplifies our derivation without loss of generality); Eq.(8) thus becomes

$$\begin{aligned}\mathcal{S} &= (+--,+++) + (+--,++-) \\ &\quad +(+--,+-+) + (+-+,+-+) \leq P_{11}. \quad (9)\end{aligned}$$

Where we have once again employed a straightforward relationship between the Wigner-symbols and the probabilities of seeing correlations at specific measurement positions. The inequality now becomes

$$P_{13} - P_{11} \leq P_{12} + P_{23}. \quad (10)$$

Note now that the inequality is written totaly in terms of measurable quantities, requiring only single outcome measurements on a bipartite system which displays some degree of anti-correlation. Assuming convex probabilities we can re-write it as

$$I_{13} - I_{\min} \leq I_{12} + I_{23}. \quad (11)$$

Where the I's indicate intensities. This constitutes the main theoretical result of this paper. Violation of this inequality negates the possibility of a local-realistic description of the world under the outlined assumptions.

Now, if we are to compare this with a system described by quantum physics, for example, an entangled pair of spin-1/2 particles, described by the pure singlet state, and taking the measurements to be positive projections onto a chosen direction given by angle θ, we have $P_{xy}(\theta_x, \theta_y) = \sin^2(\theta_x - \theta_y)$

If we make the choices $\theta_1 = 0°$, $\theta_2 = 30°$, and $\theta_3 = 60°$. Then we find: $3/4 \leq 1/4 + 1/4$. So clearly, quantum mechanics contradicts the local-realistic description of the world assumed in the formulation of the Wigner inequality. The chosen angles offer the maximal violation. For this example using pure, maximally entangled states we had $P_{11} = 0$. Our inequality allows generalizations to when this is not the case. The theoretical and experimental characterization of one such case will be the subject of the next section.

Violation with very high orbital angular momentum– A recently reported experiment on entangling states of very high orbital angular momenta of light [11] presents a difficult case if one wants to violate local realism using the generated entanglement.

For paraxial beams the total angular momentum separates out into well-defined spin [12] and orbital [13] components. Laguerre-Gauss beams naturally carry integer amounts of orbital angular momentum (OAM) in units of $\hbar$, as the quantum number l. One promising avenue of research is using the OAM degree of freedom as an infinite alphabet for quantum informatics tasks, and quantum communication [14, 15]. An important element of many such protocols is entanglement of beams with different OAM.

In our experiment, entanglement of the form $|l\rangle_a|-l\rangle_b + |-l\rangle_a|l\rangle_b$ is generated by initially creating entanglement with a Sagnac source and transferring the polarization entanglement to OAM with an interferometric scheme. For more details see Ref.[11]. The subscripts a and b label two spatial modes, and $\pm l$ is the OAM quantum number of the photon.

This state is maximally entangled in the OAM basis which spans a two-dimensional subspace of the full infinite-dimensional OAM Hilbert space. Another notable feature of this state is that its coincident-intensity, between the two beams, as a function of angular position, displays an interference pattern with $2l$ fringes. Specifically, the coincident intensity, calculated quantum mechanically, is given by $\cos^2(l(\phi_a - \phi_b))$, where ϕ_a and ϕ_b are the angular positions. Given this information, a way of determining whether or not a source is producing the target state presents itself rather intuitively. That is, placing a slit wheel with $2l$ transparent slits (the rest being opaque) in each beam, and then rotating one or the other of them so that the interference pattern can be read out by bucket detectors placed behind them as a function of slit-wheel position. If the pattern matches the expected fringes, then it can be concluded that the target state is being produced. This is illustrated in Fig.1.


$|l\rangle_a|-l\rangle_b + |-l\rangle_a|l\rangle_b$
ϕ_a
ϕ_b
coincidences
angle difference ϕ_o

FIG. 1: The setup produces two beams of entangled photons, which are then incident on slit wheels rotated to the given angles. Behind the slit wheels are bucket detectors. Fringes as a function of slit-wheel-angle difference can be seen in the coincidence counts.

This procedure though simple, intuitive, and inexpensive, is difficult to turn into a test of the classical worldview. Dichotomic measurements are not available because the slit wheel does not project in a two-dimensional space. This means inequalities of the type employed by Aspect in his third experiment [5], as well as any other dichotomic scheme, may not be employed. Furhtermore, due to the very lossy nature of each slit wheel single counts are always much greater than coincidences. Thus, any inequality which employs both coincidences and singles (such as in Ref.[10]) will be impossible to violate. However, these are not insurmountable problems for the inequality derived in the previous section, as we do not require dichotomic measurements and the problems caused by loss are less dramatic.

Before proceeding, it will be crucial to address an implication of the assumption of extended fairness. That is, that some of the noise present in the compensation term ($I_{\min}$), will be the result of the object performing the projection in the Hilbert space of the state being measured. For the case of the slit wheel a significant reduction of visibility is due to the fact that the slits are of finite width. However this effect can be predicted precisely with a theoretical model and then subtracted off from the results.

First, we write the Laguerre-Gauss beam in terms of angular position states

$$|l\rangle = \frac{1}{\sqrt{2\pi}}\int_0^{2\pi} d\phi e^{il\phi}|\phi\rangle. \tag{12}$$

The phase factor comes from the transverse phase profile of the LG modes. Now we can also write the projection operators representing the slit-wheels and bucket detectors

$$\begin{aligned}\hat{O}_a &= \sum_{n=0}^{2l-1}\int_{\frac{\pi n}{l}}^{\frac{\pi}{l}(n+W_s)} d\phi'_a|\phi'_a\rangle\langle\phi'_a|. & (13)\\ \hat{O}_b &= \sum_{n=0}^{2l-1}\int_{\frac{\pi n}{l}+\phi_o}^{\frac{\pi}{l}(n+W_s)+\phi_o} d\phi'_b \\ &\quad\times\left|\phi'_b-\frac{\pi}{2l}\right\rangle\left\langle\phi'_b-\frac{\pi}{2l}\right|. & (14)\end{aligned}$$

Where W_s is the slit width as a fraction of the distance to the next slit, and a and b label which beam the wheel is placed in. The shift in the projector states by $\pi/2l$ indicates that the data (or wheel) should be shifted one half-cycle. This relabeling causes the events to be anti-correlated as opposed to correlated. The factor ϕ_o represents the relative difference in angle between the slit wheels. We can then write the expectation value of both slit wheel operations, $\langle\hat{O}_a\hat{O}_b\rangle$, as

$$P(\phi_a,\phi_b) = W_s^2 - \frac{\cos(2l\phi_o)}{\pi^2}\sin^2(\pi W_s). \tag{15}$$

We can now make the replacement in Eq.(11)

$$I_{\min} \rightarrow I_{\max}\left(\frac{I_{\min}}{I_{\max}} - \frac{P_{\min}}{P_{\max}}\right). \tag{16}$$

Where the I's are the experimentally measured values and the P's are the predicted probabilities. This compensates for the finite slit width.

For the case of the state $|100\rangle_a|-100\rangle_b+|-100\rangle_a|100\rangle_b$ we choose the measurement angles $\phi_1 = 0°$, $\phi_2 = 0.3°$, and $\phi_3 = 0.6°$ and find $I_{13} = 5654 \pm 75$, $I_{\min} = 991 \pm 31$, $I_{\max} = 7845 \pm 89$, $I_{12} = 2202 \pm 47$, and $I_{23} = 2456 \pm 50$ in terms of total counts, assuming possionian statistics. The slit width used is $W_s = 0.149$, yielding $P_{\min} = 0.002$, and $P_{\max} = 0.043$ This results in a violation of Eq.(11) by $5026 \pm 116 \leq 4658 \pm 69$. This experiment represents the first violation of local realism with such high quanta.

Conclusions– We have derived a new version of the Wigner Inequality which can be applied to any experimental setup where some degree of anti-correlation is present between two particles. Violation of this inequality excludes the possibility of a local-realistic description of the experiment, under a set of assumptions which we have outlined. Unlike Bell and CHSH-type inequalities, the derived inequality does not require two-outcome measurements or use of singles – allowing its application to systems to which other tests of local realism can not be successfully applied. We experimentally violated the inequality in a novel system: an experiment where very high orbital angular momenta are entangled. Due to the particulars of the detection scheme, other local-realistic inequalities could not be violated.

Acknowledgements– The authors would like to thank Anton Zeilinger for initiating and motivating this work, and for suggesting that the Wigner inequality had potential in real experiments. This work was supported by the ERC Advanced Grant QIT4QAD, and the Austrian Science Fund FWF within the SFB F40 (FoQuS) and W1210-2 (CoQuS).

[1] J.S. Bell, Rev. Mod. Phys. **38**, 447 (1966).
[2] J.F. Clauser, M.A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
[3] S. J. Freedman and J. F. Clauser, Phys. Rev. Lett. **28**, 938 (1972).
[4] A. Aspect, P. Grangier, and G. Roger, Phys. Rev. Lett. **47**, 460 (1981).
[5] A. Aspect, J. Dalibard, and G. Roger, Phys. Rev. Lett. **49**, 1804 (1982).
[6] E.P. Wigner, Am. J. of Phys. **38**, 1005 (1970).
[7] S. Castelletto, I.P. Degiovanni, and M.L. Rastello, Phys. Rev. A **67**, 044303 (2003).
[8] F. A. Bovino, and I.P. Degiovanni, Phys. Rev. A **77**, 052110 (2008).
[9] M. Giustina, A. Mech, S. Ramelow, B. Wittmann, J. Kofler, J. Beyer, A. Lita, B. Calkins, T. Gerrits, S.W. Nam, R. Ursin, and A. Zeilinger, arXiv:1212.0533.
[10] J.F. Clauser, and M.A. Horne, Phys. Rev. D **10**, 526 (1974).
[11] R. Fickler, R. Lapkiewicz, W. N. Plick, M. Krenn, C. Schaeff, S. Ramelow, and A. Zeilinger, Science **338**, 640 (2012).
[12] J. Poynting, Proc. R. Soc. Lond. A Ser. A **82**, 560 (1909).
[13] L. Allen, M.W. Beijersbergen, R.J.C. Spreeuw, and J.P. Woerdman, Phys. Rev. A **45**, 8185 (1992).
[14] M. Bourennane, A. Karlsson, and Gunnar Björk, Phys. Rev. A **64**, 012306 (2001).
[15] S. Gröblacher, T. Jennewein, A. Vaziri, G. Weihs, and A. Zeilinger, New J. Phys. **8**, 75 (2006).
[16] P. Pearle, Phys. Rev. D **2**, 1418 (1970).